

and 25% to 50% of these require a thoracostomy tube. Fiberoptic bronchoscopy with transbronchial biopsy has an overall death rate of 0.1% to 0.2%. Pneumothorax occurs in 5% to 10% of patients. Brisk bleeding is uncommon. Open-lung biopsy in cases of chronic diffuse infiltrative lung disease has a mortality of 1% to 2%. This procedure requires general anesthesia.

Finally, the choice of the invasive diagnostic approach depends on the skill of the physician doing the procedure. Clearly, a lack of familiarity with the technique increases the risk to a patient and may impede the securing of diagnostic material.

THOMAS A. RAFFIN, MD

REFERENCES

- Castellino RA, Blank N: Etiologic diagnosis of focal pulmonary infection in immunocompromised patients by fluoroscopically guided percutaneous needle aspiration. *Radiology* 1979 Sep; 132:563-567.
- Gaensler EA, Carrington CB: Open biopsy for chronic diffuse infiltrative lung disease: Clinical, roentgenographic, and physiological correlations in 502 patients. *Ann Thorac Surg* 1980 Nov; 30:411-426.
- Nishio JN, Lynch JP III: Fiberoptic bronchoscopy in the immunocompromised host: The significance of a 'nonspecific' transbronchial biopsy. *Am Rev Respir Dis* 1980 Feb; 121:307-312.
- Rossiter SJ, Miller C, Churg AM, et al: Open lung biopsy in the immunosuppressed patient—Is it really beneficial? *J Thorac Cardiovasc Surg* 1979 Mar; 77:338-345.
- Wall CP, Gaensler EA, Carrington CB, et al: Comparison of transbronchial and open biopsies in chronic infiltrative lung diseases. *Am Rev Respir Dis* 1981 Mar; 123:280-285.

Management of Childhood Asthma

ASTHMA IN CHILDHOOD originates from a variety of intrinsic and acquired factors that cause bronchial hyperreactivity and impair airways clearance on a chronic, recurrent basis. Respiratory tract disease or injury in early infancy may result in this syndrome, which can persist into adulthood as recurrent bronchospasm or permanent loss of elastic recoil with chronic emphysematous changes.

Although asthma clinically exhibits a continuum of respiratory impairment, newer therapies are directed toward both the acute and chronic states of the syndrome.

Status asthmaticus is a medical emergency. The basic principles of treatment are unchanged and include correction of hypoxemia, adequate hydration and treatment of the underlying infection. While repeated epinephrine injections have been the mainstay of emergency bronchodilator therapy, more recent evidence

supports the equal efficacy of inhaled long-acting β_2 -adrenergic agents such as albuterol and fenoterol. A greater degree of bronchodilation, longer duration and less adverse reactions represent clear advantages of these agents. Continuous intravenous administration of theophylline ethylenediamine (aminophylline; dosage range, 0.65 to 0.85 mg per kg of body weight an hour, validated by serum concentrations of 10 to 20 μ g per ml), corticosteroid infusion and, on occasion, intravenously given isoproterenol have reduced the need for assisted ventilation and the mortality in childhood status asthmaticus.

Much effort has been put into the management of chronic asthma in children. Optimal management reduces impairment, promotes growth and reduces the incidence of life-threatening episodes of respiratory failure. Theophylline given by mouth remains the principal medication indicated in maintenance therapy. Blood theophylline analysis by high performance liquid chromatography (HPLC) allows titration of dosages to suit individual patients. Compliance and continual efficacy are improved with the use of sustained-release tablets and granules that are better absorbed, enhance bronchodilation and increase palatability on a 12-hour (versus 4- to 6-hour) schedule. Inhaled β_2 agents (via portable nebulizer for young children or hand-held metered-dose devices for older children) are helpful supplemental therapies. The long-acting agonists are particularly important in this context, providing sustained bronchodilation with little rebound bronchospasm. Oral β -agonists may be used but are less desirable because of the adverse effects. Short-course corticosteroid administration has been shown to abort impending status asthmaticus with little impact on adrenal function.

CHERYL D. LEW, MD

ARNOLD C. G. PLATZKER, MD

REFERENCES

- American Academy of Pediatrics, Section on Allergy and Immunology: Management of asthma. *Pediatrics* 1981 Dec; 68:874-879.
- Ben-Zvi Z, Lam C, Spohn WA, et al: An evaluation of repeated injections of epinephrine for the initial treatment of acute asthma. *Am Rev Respir Dis* 1983 Jan; 127:101-105.
- Samet JM, Tager IB, Speizer FE: The relationship between respiratory illness in childhood and chronic air-flow obstruction in adulthood. *Am Rev Respir Dis* 1983 Apr; 127:508-523.
- Tabachnik E, Scott P, Correia J, et al: Sustained-release theophylline: A significant advance in the treatment of childhood asthma. *J Pediatr* 1982 Mar; 100:489-492.

ADVISORY PANEL TO THE SECTION ON CHEST DISEASES

STANLEY ROKAW, MD

Advisory Panel Chairman

CMA Scientific Board Representative

Section Editor

Downey

GENNARO M. TISI, MD
CMA Section Chairman
University of California, San Diego

THOMAS A. RAFFIN, MD
CMA Section Secretary
Stanford University

MYRON STEIN, MD
CMA Section Assistant Secretary
Culver City

HANS EINSTEIN, MD
Immediate Past Panel Chairman
Los Angeles

RICHARD L. SHELDON, MD
Loma Linda University

GLEN LILLINGTON, MD
University of California, Davis

ARCHIE WILSON, MD
University of California, Irvine

DONALD TIERNEY, MD
University of California, Los Angeles

MICHAEL STULBARG, MD
University of California, San Francisco

RALPH C. JUNG, MD
University of Southern California

ANTONINO CATANZARO, MD
San Diego

LOWELL E. RENZ, MD
Sacramento

HOMER A. BOUSHEY, JR, MD
San Francisco